

A compound having formula (I) or any of its enantiomers of any mixture thereof, or a pharmaceutically acceptable salt thereof; wherein R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl or 2-hydroxyethyl; and R<sup>4</sup> is phenyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, heteroaryl and aryl; 3,4-methylenedioxyphenyl; benzyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, heteroaryl and aryl; heteroaryl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, heteroaryl and aryl; or naphthyl which may be substituted one or more times with substituents selected from the group consisting of halogen, CF<sub>3</sub>, CN, alkoxy, cycloalkoxy, alkyl, cycloalkyl, alkenyl, alkynyl, amino, nitro, heteroaryl and aryl. The compounds possess valuable pharmaceutical properties as monoamine neurotransmitter re-uptake inhibitors.

